

Data Sheet

BRD3 (BD1) Inhibitor Screening Assay Kit

Catalog # w42524

DESCRIPTION: *The BRD3 (BD1) Inhibitor Screening Assay Kit is designed to measure the inhibition of BRD3 bromodomain 1 (BD1) from binding to its substrate. The BRD3 (BD1) Inhibitor Screening Assay Kit comes in a convenient AlphaLISA® format, with biotinylated histone peptide substrate, assay buffer, detection buffer and purified, GST-tagged BRD3 BD1 to perform a total of 384 enzyme reactions. The key to the BRD3 (BD1) Inhibitor Screening Assay Kit is the highly specific binding of the BRD3 bromodomain 1 to the acetylated histone substrate. With this kit, only three simple steps on a microtiter plate are required. First, a sample containing BRD3 bromodomain 1 and an inhibitor of choice is incubated with the biotinylated substrate for thirty minutes. Next, acceptor beads are added, then donor beads, followed by reading the Alpha-counts.*

COMPONENTS:

Catalog #	Component	Amount	Storage	
w41043	BRD3 (29-145), BD1, GST-tag	20 µg	-80°C	(Avoid freeze/thaw cycles!)
	BET Bromodomain Ligand	400 µl	-80°C	
	Non-acetylated Ligand 1	200 µl	-80°C	
w43012	3x BRD assay buffer	4 ml	-20°C	
w43013	3x Detection buffer	3 ml	-20°C	

MATERIALS OR INSTRUMENTS REQUIRED BUT NOT SUPPLIED:

AlphaLISA® GSH acceptor beads, 5 mg/ml (PerkinElmer #AL109C)
AlphaScreen® Streptavidin-conjugated donor beads, 5 mg/ml (PerkinElmer #6760002S)
Optiplate -384 (PerkinElmer #6007290)
AlphaScreen® microplate reader

APPLICATIONS: *Useful for the study of bromodomain binding assays, screening inhibitors and selectivity profiling.*

CONTRAINDICATIONS: *Only limited amounts of DMSO can be included, as it has been shown to disrupt BRD-ligand interaction. Avoid green and blue dyes that absorb light in the AlphaScreen® signal emission range (520-620 nm), such as Trypan Blue. Avoid the use of the potent singlet oxygen quenchers such as sodium azide (NaN₃) or metal ions (Fe²⁺, Fe³⁺, Cu²⁺, Zn²⁺ and Ni²⁺). The presence of >1% RPMI 1640 culture medium leads to a signal reduction due to the presence of excess biotin and iron in this medium. MEM, which lacks these components, does not affect AlphaScreen® assays.*

STABILITY: *At least one year from date of receipt when stored as directed.*

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

REFERENCE: Muller, S., Filippakopoulos, P., Knapp, S., *Expert Rev. Mol. Med.* 2011 Sep 13;13:e29.

ASSAY PROTOCOL:

All samples and controls should be tested in duplicate. Use slow shaking for all incubations.

Step 1:

- 1) Prepare the master mixture: $N \text{ wells} \times (2.5 \mu\text{l } 3\times \text{ BRD assay buffer} + 1 \mu\text{l BET Bromodomain Ligand} + 1.5 \mu\text{l H}_2\text{O})$.
- 2) Thaw **BRD3 (BD1)** on ice. Upon first thaw, briefly spin tube containing protein to recover full content of the tube. Aliquot both proteins into single use aliquots. Store remaining undiluted protein in aliquots at -80°C immediately. Note: **BRD3** is very sensitive to freeze/thaw cycles. Do not re-use thawed aliquots or diluted protein.
- 3) Dilute **BRD3 (BD1)** in **1x BRD assay buffer** at $1.6 \text{ ng}/\mu\text{l}$. Keep diluted proteins on ice until use. Discard any unused diluted protein after use.

Add $5 \mu\text{l}$ of master mixture to each well designated for the "Positive Control", "Test Inhibitor", and "Blank". For the "Substrate Control", add $2.5 \mu\text{l } 3\times \text{ BRD assay buffer} + 1 \mu\text{l Non-acetylated Ligand 1} + 1.5 \mu\text{l H}_2\text{O}$.

	Blank	Substrate Control	Positive Control	Test Inhibitor
3x BRD assay buffer	$2.5 \mu\text{l}$	$2.5 \mu\text{l}$	$2.5 \mu\text{l}$	$2.5 \mu\text{l}$
BET Bromodomain Ligand	$1 \mu\text{l}$	–	$1 \mu\text{l}$	$1 \mu\text{l}$
Non-acetylated Ligand 1	–	$1 \mu\text{l}$	–	–
H ₂ O	$1.5 \mu\text{l}$	$1.5 \mu\text{l}$	$1.5 \mu\text{l}$	$1.5 \mu\text{l}$
Test Inhibitor/Activator	–	–	–	$2.5 \mu\text{l}$
Inhibitor buffer (no inhibitor)	$2.5 \mu\text{l}$	$2.5 \mu\text{l}$	$2.5 \mu\text{l}$	–
1x BRD assay buffer	$2.5 \mu\text{l}$			
BRD3 (BD1) ($1.6 \text{ ng}/\mu\text{l}$)	–	$2.5 \mu\text{l}$	$2.5 \mu\text{l}$	$2.5 \mu\text{l}$
Total	$10 \mu\text{l}$	$10 \mu\text{l}$	$10 \mu\text{l}$	$10 \mu\text{l}$

- 4) Add $2.5 \mu\text{l}$ of inhibitor solution to each well designated "Test Inhibitor". For the "Positive Control", "Substrate Control" and "Blank", add $2.5 \mu\text{l}$ of the same solution without inhibitor (inhibitor buffer). Note: Keep DMSO concentration below 0.5 %.
- 5) Add $2.5 \mu\text{l}$ of **1x BRD assay buffer** to the well designated "Blank".

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

- 6) Initiate reaction by adding 2.5 μ l of diluted **BRD3 (BD1)** prepared as described above to each well labeled "Positive Control", "Test Inhibitor", and "Substrate Control". Incubate at room temperature for 30 minutes.

Step 2:

Note: Protect your samples from direct exposure to light!


- 1) Dilute GSH Acceptor beads (PerkinElmer #AL109C) 250-fold with 1x Detection buffer. Add 10 μ l per well. Shake plate briefly. Incubate at room temperature for 30 minutes.

Step 3:

- 1) Dilute Streptavidin-conjugated donor beads (PE #6760002S) 250-fold with 1x Detection buffer. Add 10 μ l per well. Incubate at room temperature for 15 – 30 minutes.
- 2) Read Alpha-counts.

Due to lot to lot variability in AlphaScreen® bead performance, it may be necessary to optimize assay conditions. For example, slight adjustments to bromodomain or ligand concentrations may improve signal-to-noise ratio.

Example of Assay Results:



BRD3 (BD1) binding activity, measured using the BRD3 (BD1) Inhibitor Screening Assay Kit, West Bioscience, Catalog #w42524 and (+)-JQ1 Inhibitor, Catalog #w37411. Data shown is lot-specific. For lot-specific information, please contact West Bioscience, Inc. at sale@westbioscience.com.

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



9891 Irvine Center Dr. Suite 200

Irvine, CA 92618

United States

Tel: 1.800.831.1518

Fax: 1.800.831.1518

Email: sale@westbioscience.com

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.westbioscience.com



9891 Irvine Center Dr. Suite 200

Irvine, CA 92618

United States

Tel: 1.800.831.1518

Fax: 1.800.831.1518

Email: sale@westbioscience.com

OUR PRODUCTS ARE FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

www.westbioscience.com